

530536

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 April 2004 (22.04.2004)

PCT

(10) International Publication Number
WO 2004/032655 A1

(51) International Patent Classification⁷: A23L 3/015,
3/3571

(74) Agents: CALHOUN, Douglas, C., et al.; A J Park, 6th
Floor Huddart Parker Building, Post Office Square, P.O.
Box 949, 6015 Wellington (NZ).

(21) International Application Number:
PCT/NZ2003/000224

(22) International Filing Date: 8 October 2003 (08.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
521836 8 October 2002 (08.10.2002) NZ

(71) Applicant (*for all designated States except US*):
FONTERRA CO-OPERATIVE GROUP LIMITED
[NZ/NZ]; Building 103, Leonard Isitt Drive, Auckland
Airport, Auckland (NZ).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **CARROLL, Tim**
[IE/NZ]; 51 Collingwood Street, Palmerston North
(NZ). **CHEN, Ping** [NZ/CN]; 10-2006, JiaheYuan, 148
Donfengxi Road, Liwan, Guangzhou (CN). **HARNETT,**
Michelle [NZ/NZ]; 2 Anuru Place, Palmerston North (NZ).
HARNETT, James [NZ/NZ]; 237 Aokautere-Pahiatua
Road, RDI, Palmerston North (NZ).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PRESSURE TREATING FOOD TO REDUCE SPOILAGE

(57) Abstract: The present invention broadly describes a method of pressure treating foods containing cultures, wherein the pressure treatments are performed under such conditions such that cultures survive while the growth of spoilage microflora is reduced, delayed, prevented or eliminated. Foods treated according to the invention include cultural dairy foods (such as yoghurts), fruit and vegetable juices and other dairy foods (such as cheese).



WO 2004/032655 A1

PRESSURE TREATING FOOD TO REDUCE SPOILAGE

FIELD OF INVENTION

5 The present invention relates to methods of pressure treating food so as to reduce spoilage of the food. More particularly the invention relates to the use of high pressure treatment to reduce microbial spoilage in foods and / or to render the food safe for consumption, while retaining viable desired cultures.

10 BACKGROUND

Many foods have relatively short shelf lives due to the presence of undesired contaminants such as yeasts and moulds. Such yeasts and moulds cause undesired spoilage, and often render the foods inedible.

15

It is known to inactivate undesirable microorganisms in food by a variety of methods, the most common of which is heating. Heat treatment can significantly improve both the safety and the keeping quality of the food. In particular the shelf life of the food can be extended.

20

However, properties such as the taste, texture, and nutritional quality of some foods can be compromised by a heat treatment. For example, heat-treated meat can have unacceptable cooked flavour. A heat-treated cultured milk product such as yoghurt does not contain a live bacterial culture, as the culture is inactivated by the treatment.

25

It was recognised at the beginning of the last century, that bacteria used for the fermentation of products such as yoghurt, are of benefit to human health if consumed alive. It is now recognised that certain cultures of live micro-organisms, defined as probiotic, exert health benefits beyond basic nutrition upon ingestion in certain numbers.

30

(Holzapfel et al). It is known to add these probiotic bacteria to foods (Lee and Salminen)

for delivery by ingestion. However, it is difficult to deliver such bacteria in sufficient numbers in a food that is subsequently heat-treated.

- 5 It is an object of the present invention to provide an improved or alternative method of treating a food product, and / or to go at least some way to overcoming the problems encountered with the prior art.

SUMMARY OF INVENTION

10

In one aspect the invention broadly comprises of a method of treating a food comprising the following steps:

- selecting a food comprising at least one strain of a culture, said strain capable of surviving a pressure treatment at a predetermined pressure and pH, and
- 15 - subjecting the food to a treatment pressure at or below the predetermined pressure, wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora.

20 Useful treatment pressures according to the invention may be selected from 350MPa, 360MPa, 370MPa, 380MPa, 390MPa, 400MPa, 410MPa, 420MPa, 430MPa, 440MPa, 450MPa, 460MPa, 470MPa, 480MPa, 490MPa, 500MPa, 510MPa, 520MPa, 530MPa, 540MPa, 550MPa, 560MPa, 570MPa, 580MPa, 590MPa, 600MPa, 610MPa, 620MPa, 630MPa, 640MPa and 650MPa.

25

Preferably the food is subjected to a pressure of at least 350MPa.

More preferably the food is subjected to a pressure of at least 400MPa.

30 It is envisaged that the invention may be performed at a pH level selected from the following: 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7,

4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8; 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9 and 8.0.

5 In the preferred embodiment, the food is at a pH of between 3.0 and 8.0 when subjected to the treatment pressure.

Preferably the pH is between 3.6 and 4.8.

Most preferably the pH is between 4.0 and 4.6.

10

Preferred temperatures at which the invention may be performed can be selected from (in degrees Celsius): 0, 4, 5, 10, 15, 20, 25, 30, 35 and 40.

15 Preferably the food is pressure treated at a temperature in the range 0 degrees Celsius to 40 degrees Celsius.

Most preferably the food is pressure treated at a temperature in the range 0 degrees Celsius to 20 degrees Celsius.

20 In a preferred embodiment, the food is pressure treated after packaging.

A preferred food according to this aspect of the invention is a cultured dairy product.

A preferred cultured dairy product used in the invention is yoghurt.

25

Alternative cultured dairy products used in the invention may be selected from yoghurt drinks, dairy desserts, cottage cheese, cream cheese and cultured beverages.

Preferred strains of culture used in this aspect of the invention are selected from:
30 *Lactobacillus acidophilus* HN017 AGAL deposit number NM97/09515 dated 18 August 1997, *Bifidobacterium lactis* HN019 AGAL deposit number NM97/09513 dated 18

August 1997, *Streptococcus thermophilus* St10, *Streptococcus thermophilus* St49, *Lactobacillus helveticus* Lh1, *Lactobacillus helveticus* Lh5001, *Lactobacillus delbrukeii subsp bulgaricus* Lb1, Rhodia MY900 (commercially sold by Rhodia under the trade mark "MY900"), Rhodia MY105, Rhodia MYE95, Rhodia MYBio6, Rhodia TA060, Rhodia LH100, Chr. Hansen ABT4, Chr. Hansen YC-X11, Chr. Hansen ABT3, Danisco V1, Danisco Yo Mix VW, Danisco MSK Mix ABN1-45, *Bifidobacterium lactis* Bb12 (commercially sold by Nestle under the trade mark "Bb12"), *Bifidobacterium lactis* Wisby 420 (commercially sold by Wisby under the trade mark "420") and combinations thereof. The strains identified as St10, St49, Lh1, Lh5001 and Lb1 are commercially available from the Fonterra Research Centre Limited, Palmerston North, New Zealand.

In a second aspect, the invention broadly comprises of a method of treating a food, comprising the steps:

- selecting a food containing at least one strain of a culture, said strain being a probiotic strain capable of surviving a pressure treatment at a predetermined pressure and pH, and
- subjecting the food to a treatment pressure at or below the predetermined pressure, wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora.

It is envisaged that the probiotic may be either used to ferment the food, or may be added to the food directly.

Probiotic strains used in the invention may be selected from strains of *Bifidobacterium*, preferably *Bifidobacterium lactis*.

Preferred probiotic strains used in the invention are selected from *Bifidobacterium lactis* HN019 AGAL deposit number NM 97/09513 dated 18 August 1997, and *Bifidobacterium* sold under the trade names Bb12 (Nestle) and Wisby 420.

Other preferred probiotic strains used in the invention are selected from strains of *Lactobacillus*, preferably *Lactobacillus acidophilus*.

Most preferably a probiotic strain used in the invention is *Lactobacillus acidophilus*
5 HN017 AGAL deposit number NM 97/09515 dated 18 August 1997.

Useful treatment pressures according to the invention may be selected from 350MPa, 360MPa, 370MPa, 380MPa, 390MPa, 400MPa, 410MPa, 420MPa, 430MPa, 440MPa, 450MPa, 460MPa, 470MPa, 480MPa, 490MPa, 500MPa, 510MPa, 520MPa, 530MPa,
10 540MPa, 550MPa, 560MPa, 570MPa, 580MPa, 590MPa, 600MPa, 610MPa, 620MPa, 630MPa, 640MPa and 650MPa.

More preferably the pressure is at least 350MPa.

15 Even more preferably the pressure is at least 400MPa.

Alternatively the pressure is at least 500MPa.

It is envisaged that the invention may be performed at a pH level selected from the
20 following: 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5 and 4.6.

In the preferred embodiment, the food is at a pH of between 3.0 and 4.6 when subjected to the treatment pressure.

25 Preferred conditions of temperature are as noted for the first aspect of the invention.

In a third aspect the invention broadly comprises of a method of treating a food comprising the following steps:

- selecting a food containing at least one strain of a protective culture, said
30 strain capable of surviving a pressure treatment at a predetermined pressure and pH, and

- subjecting the food to a treatment pressure at or below the predetermined pressure, wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora.
- 5 Preferably the protective culture is selected from those used in cultured dairy foods, fermented foods, cooked meats, vegetables, salads, cook-chilled foods, ready-to-eat foods. Such protective cultures include, but are not limited to, probiotics, bacteriocins and acid producing bacteria.
- 10 Preferred conditions of pH and temperature are as noted for the first aspect of the invention.

In a fourth aspect the invention consists in the use of at least one bacterial strain in a food to be subjected to a pressure treatment at a predetermined pressure such that undesired

15 microflora are inactivated while the bacterial strain survives, said bacterial strain being selected from: *Lactobacillus acidophilus*, *Bifidobacterium lactis*, *Lactobacillus acidophilus* HN017 AGAL deposit number NM97/09515 dated 18 August 1997, *Bifidobacterium lactis* HN019 AGAL deposit number NM97/09513 dated 18 August 1997, *Streptococcus thermophilus* St10, *Streptococcus thermophilus* St49, *Lactobacillus*

20 *helveticus* Lh1, *Lactobacillus helveticus* Lh5001, *Lactobacillus delbrueckii subsp bulgaricus* Lb1, Rhodia MY900, Rhodia MY105, Rhodia MYE95, Rhodia MYBio6, Rhodia TA060, Rhodia LH100, Chr. Hansen ABT4, Chr. Hansen YC-X11, Chr. Hansen ABT3, Danisco V1, Danisco Yo Mix VW, Danisco MSK Mix ABN1-45, and *Bifidobacterium* sold under the trade names Bb12 (Nestle) and Wisby 420 (Wisby).

25

According to the aspects of the invention, the foods may be subjected to the treatment pressure for between about 1 second and about 10 minutes. Preferred times may be selected from 1 second, 5 seconds, 10 seconds, 20 seconds, 30 seconds, 60 seconds, 90 seconds, 2 minutes, 3 minutes, 4 minutes, 5 minutes, 6 minutes, 7 minutes, 8 minutes, 9

30 minutes or 10 minutes.

The invention also consists in a food treated by the methods described herein.

The invention may also broadly be said to consist in any alternative combination of features as described or shown in the accompanying examples. Known equivalents of these features not expressly set out are nevertheless deemed to be included.

BRIEF DESCRIPTION OF FIGURES

Figures 1a and 1b are graphs showing the effect of varying pressure treatments on Penicillium mould and yoghurt starter (MY900 – sold by Rhodia).

Figure 2 shows the effect of varying pressure treatments on levels of viable culture and spoilage organisms after 95 days of storage at 4 degrees Celsius.

Figure 3 shows the effect of varying pressure treatments on levels of viable culture and spoilage organisms after 15 days of storage at 20 degrees Celsius.

Figure 4 shows the effect of pressure treatments performed at different pH levels.

DETAILED DESCRIPTION

As mentioned herein, references to “pressure treatment” or “UHP treatment” mean ultra high-pressure treatments. Such treatments are generally accepted as pressure treatments using pressures of at least 100MPa. This is also known in the art as “high pressure”, “high hydrostatic pressure” (HHP) or “high pressure processing” (HPP).

A pressure treatment is understood to comprise the following steps:

- placing a food into the chamber and sealing the chamber,
- raising the pressure in the chamber, and thereby the food to a predetermined set pressure,

- holding the food at this pressure for a predetermined time (termed the treatment time, dwell time or hold time), and
- releasing the pressure from the chamber and removing the food.

5 The characteristics of the high-pressure equipment used might affect the conditions required to successfully perform the invention. In particular, the time taken to achieve the treatment pressure and to release the treatment pressure from the food, and the accuracy with which the treatment pressure is delivered and controlled may influence the outcomes, particularly in situations where it is not necessary for the food to be held at the
10 treatment pressure for an appreciable time.

The nature of the pressure treatment process results in temperature fluctuations in the treated food during treatment. As such, references to preferred temperatures during pressure treatments refer to the temperature of the food or beverage before the pressure is
15 raised.

As mentioned herein, references to "food" or "foodstuffs" include, for example yoghurts, yoghurt drink, Kefir, cheeses, milk, dairy products, dairy desserts, fruit juices, beverages, sport drinks, and the like.

20

As mentioned herein, references to "spoilage microflora" refer to contaminants such as yeasts and moulds, food-poisoning bacteria, pathogens, naturally present bacteria, and starter organisms that have completed their function. References to "spoilage" refer to the presence of such organisms in food or foodstuffs, and the effect that this presence has on
25 the various properties of the food (e.g. shelf life, taste or texture).

References are made to the reduction, delay, prevention and elimination of spoilage microflora. Such references encompass situations where existing spoilage microflora are inactivated, in particular where such inactivation is required to meet food safety labeling
30 or regulatory requirements.

As mentioned herein, references to “protective culture” refer to a live culture that produces a metabolite that exhibits antimicrobial activity.

5 As mentioned herein, references to “probiotics” refer to strains of bacteria with health promoting and immune-enhancing properties. Such strains have an ability to survive in and colonise the intestine and gut, and are well known in the art.

10 As mentioned herein, references to “surviving” a pressure treatment refer to the survival of strains in useful numbers. For example, in yoghurts, “surviving” a pressure treatment means that the number of intact viable starter organisms after the pressure treatment are such that they satisfy standards of identity, product definition and regulation (typically one hundred thousand colony forming units per gram or higher). With reference to probiotics, “surviving” may refer to the number of intact probiotic organisms required for a dosage when ingested with a quantity of food (typically one hundred thousand colony
15 forming units per gram or higher).

A desirable feature of UHP treatment is its non-invasive nature. With appropriate equipment, foodstuffs and the like can be treated in the same containers that they are to be distributed to the consumer in. For example, set or stirred yoghurt can be treated in
20 pottles and cultured dairy drinks can be treated in their bottles. Pressure treatment may also be applied to non-homogeneous foods such as yoghurts containing fruit, nuts or fruit purees.

It is also envisaged that foods can be treated in accordance with the invention when they
25 have already contain appreciable numbers of spoilage microflora, and are considered “borderline” by consumer and / or regulatory requirements.

For situations where the desired microorganisms (for example, yoghurt cultures and / or probiotics) and the undesired microorganisms such as contaminants are in the same
30 sealed container, a method of selective inactivation has been identified.

It has now been identified that it is possible to subject a food to a UHP treatment that inactivates spoilage microflora or other unwanted microorganisms, but in which desirable bacteria survive the treatment in useful numbers. These desirable bacteria could be typical yoghurt or probiotic cultures that offer a health benefit if consumed alive or live protective cultures that can inhibit contaminating microorganisms.

The invention offers the ability to preserve the quality of foods containing a live culture that is of significant commercial value. In particular we have found that we can produce cultured dairy products (such as yoghurt that contains abundant live cultures) having an extended shelf life due to reduced spoilage by spoilage microflora, such as yeasts and moulds.

The invention may also be used to produce a range of food products containing an abundant live probiotic culture, with an improved keeping quality due to reduced microbial contamination or spoilage.

Also identified is a method of treating foods containing a live protective culture that is capable of inactivating spoilage microflora. This process can be used to improve the safety and keeping quality of certain foods.

One method of identifying appropriate cultures for use in the present invention is to inoculate strains into a food and then treat the food at a pressure suitable for controlling the undesired microorganism, for example 350 MPa for 5 minutes or 450 MPa for 1 minute or 460 MPa for 1 second or 600 MPa for 1 second. Strains that survive the test-pressure treatment in useful numbers are identified as strains that may be useful according to the present invention.

In the case of yogurts, "useful numbers" can mean the number of viable starter (or specified microorganisms) specified by standards of identity, product definitions or regulation, (typically one hundred thousand colony forming units per gram or higher). In the case of foods containing viable probiotic bacteria, useful numbers can mean the

number of viable bacteria required such that a sufficient dose is ingested with a specified quantity of the food (typically one hundred thousand colony forming units per gram or higher).

- 5 Once the suitably tolerant strains of culture have been identified, a food product can be manufactured containing that strain of culture. Alternatively, food products known to contain pressure tolerant bacteria could be identified. The food products can then be subjected to pressure treatment under conditions which are known to inactivate spoilage microflora, but allow survival of the selected strains of culture.

10

The method of selective inactivation can be understood with reference to Figures 1a and 1b which compares the levels of desired bacteria (commercial starter Rhodia MY900) and spoilage microflora (an intentionally added *Penicillium* mould) in a yoghurt at pH 4.4. In Figure 1a, the yoghurt is subjected to various treatment pressures for a time of one
15 second. After pressure treatment at about 460 MPa, the number of viable starter bacteria is one hundred million colony-forming units per gram or higher, but the mould is inactivated to levels below detection limits. After a pressure treatment of about 430 MPa, the number of viable starter bacteria is one hundred million colony-forming units per gram or higher, but the mould is reduced by at least 2.4 log cycles in number (this may be
20 sufficient to prevent spoilage in a yogurt with a pre-treatment quality specification of less than one hundred colony-forming units of mould per gram). After a pressure treatment of less than 410 MPa, there is increased presence of mould. In Figure 1b, the yoghurt is subjected to various treatment pressures for a time of five minutes. After a pressure treatment of about 400 MPa, the number of viable starter bacteria is one hundred million
25 colony-forming units per gram or higher. After a pressure treatment of about 350 MPa the mould numbers are reduced by at least one log cycle, and at a pressure of about 400 MPa, the mould numbers are reduced to levels below detection limits. However at a pressure of 450 MPa or higher, although the mould numbers are reduced to levels below the limits of detection, the number of viable starter bacteria declines appreciably from about one
30 million per gram at 500 MPa to only about 10 per gram at 600 MPa. Pressure treatment conditions may be selected based on considerations such as the level of protection against

mould spoilage required, and the level of contamination before treatment as well as the number of viable desirable bacteria required.)

Figures 1a and 1b show the how varying pressure treatments can be used to inactivate an undesired microorganism (in this instance *Penicillium* mould shown on the graph as solid symbols) while retaining a desired microorganism in useful numbers (*Rhodia* MY900 yoghurt starter – open symbols). The grey highlighted portions illustrate a treatment window within which the competing requirements of desirable microorganism viability and undesired contaminant inactivation may be satisfied.

Some cultures can produce anti-microbial metabolites (such as bacteriocins), that impart specific protection against certain microorganisms (Holzapfel et al, Caplice & Fitzgerald). Such protective cultures may be added directly to the food as a viable starter organism or starter adjunct organism to improve microbial safety of the food and particularly protect against temperature abuse on distribution and storage. An alternative way to use such cultures is to first ferment an appropriate substrate to produce the desired metabolites and then inactivate the culture for sale and subsequent incorporation into the food.

Probiotic bacteria can be considered as an example of a protective culture by virtue of their influence on resident intestinal flora (Holzapfel et al Lee & Salamin).

Protective cultures can be used in cultured dairy foods, fermented foods, cooked meats, vegetables, salads, cook-chill foods and ready-to-eat meals.

Using the methods taught in the present invention, it is now possible to extend the shelf life and / or reduced spoilage levels of some foods by inactivating spoilage microflora such as spoilage organisms whilst retaining the viable protective cultures in the food to protect against growth of other undesired microflora, such as bacteria or bacteria spores.

The invention consists in the foregoing and also envisages constructions of which the following gives examples.

EXAMPLES

5

In some of the following examples, typical contaminants were deliberately added to the food products. The food products were then treated according to the methods of the invention.

- 10 The methods used to enumerate the bacteria in the following examples are summarised in Table 5 ("cfu" = colony forming units). All cultures were obtained from the Fonterra Research Centre Culture Collection ("FRCCC") in Palmerston North, New Zealand, unless otherwise stated. The contaminant cultures used in the examples and their respective origins are summarised in Table 6.

15

The pressure treatments were done on Stansted Fluid Power FoodLab laboratory (30 mL) and pilot (675mL) vessels (except for Examples 10 and 11, done on a Flow Autoclave Systems 2L unit).

- 20 In examples where a treatment temperature is given, this was the temperature of the chamber contents before the pressure was raised.

Example 1: A cultured dairy product containing a viable culture

- 25 A 10% reconstituted skim milk (RSM) substrate was inoculated with 1% of a culture of *Streptococcus thermophilus* St-10 (FRCCC) and fermented overnight at 37°C. The pH of the cultured skim milk was adjusted to 4.4 by addition of lactic acid, and intentionally contaminated by spiking at 1.8×10^6 cfu/g (Table 5) with a *Yarrowia lipolytica* yeast (Table 6). The contaminated cultured milk so produced was then treated by applying a
- 30 pressure of 400 MPa for 5 minutes at 10°C. This process produced a product with no

detectable contaminating yeast (>5-log inactivation), while retaining a starter culture count of 3.3×10^7 cfu/g (M17 agar, Table 5).

Example 2 : A cultured dairy product containing a viable culture

5

A 10% RSM substrate was inoculated with 1% of a culture of *Lactobacillus helveticus* Lh-5001 (FRCCC) and fermented overnight at 37°C. The pH of the cultured skim milk was adjusted to 4.4 and intentionally contaminated by spiking to 3.1×10^6 cfu/g (Table 5) with a *Penicillium* mould (Table 6). The contaminated cultured milk so produced was then treated by applying a pressure of 400 MPa for 5 minutes at 10°C. This process produced a product with no detectable contaminating mould, while retaining a starter culture count of 1.3×10^8 cfu/g (MRS agar, Table 5).

Example 3 : A cultured dairy product containing a viable culture

15

A 10% RSM substrate was inoculated with 1% of a culture of *Lb. delbruekii sub-species bulgaricus* Lb-1 (FRCCC) and fermented overnight at 37°C. The pH of the cultured skim milk was adjusted to 4.4, and intentionally contaminated by spiking at 9.3×10^7 cfu/g (Table 5) with a *Debromyces hanseii* yeast (Table 6). The contaminated cultured milk so produced was then treated by applying a pressure of 350 MPa for 5 minutes at 10°C. This process produced a product with no detectable contaminating yeast, while retaining a starter culture count of 7.6×10^7 cfu/g (MRS agar, Table 5).

Example 4 : Cultured dairy product with no significant live viable culture [control example]

25

A 10% RSM substrate was inoculated with 1% of a culture of *S. thermophilus* St-1 (FRCCC) and fermented overnight at 37°C. The pH of the cultured skim milk was adjusted to 4.4 by addition of lactic acid, and intentionally contaminated by spiking at 3.5×10^6 cfu/g (Table 5) with a pink yeast (Table 6). The contaminated cultured milk was then treated by applying a pressure of 400 MPa for 5 minutes at 10°C. This process

30

produced a product with no detectable contaminating yeast (>5-log inactivation). However, the starter culture was reduced from 4.4×10^7 cfu/g to 6.8×10^2 cfu/g (>4-log inactivation) by the pressure treatment.

5 Example 5 : A yoghurt containing a viable culture

A yoghurt milk made up with 7.0% skim milk powder (SMP) and 7.5% whole milk powder (WMP), was heated to 55°C and homogenised at 150/50 bar. The homogenised milk was then heated to 90°C in a steam-heated waterbath and held at that temperature for 10 minutes. After rapid cooling to 42°C, the milk was inoculated with 1% of a culture of *S. thermophilus* St-10 and *Lb. delbruekii sub-species bulgaricus* strain Lb-5033 (FRCCC), then fermented at 42°C to a pH of 4.4, whereupon it was cooled to 4°C. The resulting yoghurt was intentionally contaminated by spiking at 4.4×10^6 cfu/g with a *Penicillium* mould and pink yeast. The contaminated yoghurt was then treated by applying a pressure of 450 MPa for about 1 minute at 15°C. This process produced a product with no detectable contaminating yeast or mould (>5-log inactivation), while retaining a starter culture count of 3.0×10^8 cfu/g (M17 agar, Table 5) and 1.4×10^8 cfu/g (MRS agar, Table 5).

20 Example 6 : A fruit yoghurt containing a viable culture

7% of a sterile fruit puree was added to yoghurt made as in Example 5. It was intentionally contaminated by spiking at 5.4×10^6 cfu/g with a *Penicillium* mould and pink yeast. The contaminated yoghurt was then treated by applying a pressure of 450 MPa for about 1 minute at 15°C. This process produced a product with no detectable contaminating yeast or mould (>5-log inactivation), while retaining a starter culture count of 6.6×10^8 cfu/g (M17 agar, Table 5).

Example 7 : A yoghurt drink containing a viable culture

- 5 The yoghurt, produced as in Example 5, was made up to a final formulation of 8% sugar, 1% protein and 0.4 % carboxymethylcellulose .The pH of the yoghurt was then adjusted to 4.0 by addition of a citric/lactic acid solution and homogenised at 200 bar. The resulting yoghurt drink was intentionally contaminated by spiking at 7.9×10^6 cfu/mL with a *Penicillium* mould and pink yeast. The contaminated yoghurt drink was then treated by
- 10 applying a pressure of 450 MPa for about 1 minute at 15°C. This process produced a product with no detectable contaminating yeast (>5-log inactivation), while retaining a starter culture count of 2.1×10^8 cfu/mL (M17 agar, Table 5).

Example 8 : A yoghurt containing a viable culture

- 15 A yoghurt milk made up with 7.0% SMP and 7.5% WMP, was heated to 55°C and homogenised at 150/50 bar. The homogenised milks were then heated to 90°C in a steam-heated waterbath and held at that temperature for 10 minutes. After rapid cooling to 42°C, the yoghurt milk was inoculated with 1% each of *S. thermophilus* St-10 and *Lb. delbruekii sub-species bulgaricus* strain Lb-1 and fermented at 42°C to a pH of 4.0,
- 20 whereupon it was cooled to 4°C. The yoghurt was contaminated by spiking at 3.0×10^6 cfu/g with a *Penicillium* mould and a pink yeast, and treated by applying a pressure of 450 MPa for about 1 minute at 15°C. This process produced a product with no detectable contaminants (>5-log inactivation), while retaining a starter culture count of 1.8×10^8
- 25 cfu/g (M17 agar, Table 5) and 4.1×10^7 cfu/g (MRS agar, Table 5).

Example 9 : A yoghurt with no significant viable culture after treatment at a low final pH

5

A yoghurt milk made up with 7.0% SMP and 7.5%WMP, was heated to 55°C and homogenised at 150/50 bar. The homogenised milks were then heated to 90°C in a steam-heated waterbath and held at that temperature for 10 minutes. After rapid cooling to 42°C, the yoghurt milk was inoculated with 1% each of *S. thermophilus* St-10 and *Lb. delbruekii sub-species bulgaricus* strain Lb-1 and fermented at 42°C to a pH of 3.6, whereupon it was cooled to 4°C. The yoghurt was intentionally contaminated by spiking at 3.0×10^6 cfu/g with a *Penicillium* mould and a pink yeast, and treated by applying a pressure of 450 MPa for about 1 minute at 15°C. This process produced a product with no detectable contaminants (>5-log inactivation). However, the starter culture count was reduced to 1.3×10^4 cfu/g, from 3.2×10^8 cfu/g (M17 agar, Table 5).

15

Example 10: A yoghurt containing a live probiotic culture

A yoghurt milks made up with 7.0% SMP and 7.5% WMP, was heated to 55°C and homogenised at 150/50 bar. The homogenised milks were then heated to 90°C in a steam-heated water bath and held at that temperature for 10 minutes. After rapid cooling to 42°C, the yoghurt milk was inoculated with 1% *B. lactis* HN019, 0.25% ST10 and 0.25% Lb TH (*Lb. delbruekii sub-species bulgaricus* TH, FRCCC). The yoghurt milk was incubated at 38°C overnight (16 h) to a pH of 4.0. removed from the incubator and cooled to approximately 25°C. It was then smoothed by passing it through a homogeniser at 0 bar and filled into 375g PET bottles and stored at 5°C. The yoghurts were pressure-treated at 375 MPa for 5 minutes.

20

25

An untreated yoghurt sample was retained as a control.

30

After four weeks storage at 4°C, the yoghurts contained a live probiotic culture (*B. lactis* HN019) of 1.3×10^7 cfu/g (RBA agar, Table 5), with no contaminating yeasts and moulds.

- 5 In contrast, although the untreated control sample contained a live probiotic culture of 2.0×10^7 cfu/g, it was contaminated with 1.2×10^3 cfu/g of yeasts and moulds after 4 weeks at 4°C.

Example 11: A yoghurt containing a live probiotic culture

10

A yoghurt formulated as 7.0% SMP, 7.5%WMP, 0.1% pectin and 0.4% starch was prepared as for Example 10. The yoghurt was pressure-treated at 375 MPa for 5 minutes, and untreated yoghurt was retained as a control. After the treated sample was stored at 4°C for 4 weeks, it contained a live probiotic culture of 5.6×10^7 cfu/mL (RBA agar, 15 Table 5), with no contaminating yeasts and moulds. In contrast, although the untreated control sample contained a live probiotic culture of 6.2×10^7 cfu/mL(after 4 weeks at 4°C) it was contaminated with 3.2×10^2 cfu/mL of yeasts and moulds.

Example 12: A direct acidified milk drink containing a live probiotic culture

20

A direct acidified milk drink was made by blending Carboxymethylcellulose (CMC) with sugar, and then dispersing it in 1.5 L warm (55°C) water. WMP was separately dispersed in 1.5 L warm (55°C) water and then mixed with the CMC-sugar solution. More sugar was added to give a final formulation of 8% sugar, 3.2% WMP and 0.4 % CMC. The pH 25 of the milk drink was then adjusted to 4.0 by addition of a citric/lactic acid solution, homogenised at 200 bar, and sterilised at 85°C in steam bath for 5 minutes. The drink was then chilled and inoculated with 5.7×10^7 cfu/mL (RCA agar, Table 5) of *Bifidobacterium* (Wisby strain 420). The probiotic drink produced was then treated by applying a pressure treatment of 350 MPa for 5 minutes at room temperature (15 °C). A 30 drink containing live probiotic *Bifidobacterium* at the level of 4.7×10^7 cfu/mL (RCA agar, Table 5) was so produced.

Example 13: A direct acidified milk drink containing no significant live probiotic culture [control example]

- 5 The direct-acidified milk drink made as in Example 12 was inoculated with at 4.9×10^7 cfu/mL (MRS agar, Table 5) of *L. casei* Yakult Shirota strain. The sample was treated at a pressure of 350 MPa for 5 minutes at 15 °C. This process produced a product with a count of only 240 cfu/mL (5.3-log reduction) of the *L. casei* probiotic culture after the pressure treatment.

10

Example 14: An orange juice containing a live probiotic culture

- Australian navel oranges were squeezed to produce an orange juice of pH 3.48 and total solids of 10.6 %. *B. lactis* strain HN019 was inoculated into the juice at 1.1×10^8 cfu/mL (RCA agar, Table 5) and the juice was intentionally contaminated by spiking at 2.2×10^7 cfu/mL with pink yeast. The contaminated probiotic orange juice was then treated at a pressure of 350 MPa for 5 min at 15 °C. This process produced a product with no detectable contaminating pink yeast (> 6-log inactivation) while retaining a probiotic culture count of 1.1×10^8 cfu/mL (RCA agar, Table 5).

20

Example 15: An orange juice containing a live probiotic culture and undesirable organisms

- The contaminated orange juice made as in Example 14 was treated at a pressure of 300 MPa for 5 minutes at 15°C. Although, the product retained a probiotic culture count of 1.1×10^8 cfu/mL (RCA agar, Table 5), it was contaminated by 3.8×10^3 cfu/mL of pink yeast.

25

Example 16: An orange juice containing a live probiotic culture

The contaminated orange juice made as in Example 14 was treated at a pressure of 600 MPa for 5 minutes at 15 °C. This process produced a product with no detectable contaminating Pink yeast (> 6-log inactivation) while retaining a probiotic culture count of 4.1×10^6 cfu/mL (RCA agar, Table 5).

Example 17: Method of strain selection

10

Potential probiotic strains were screened for suitability in the production of a pressure-treated direct-acidified milk drink as follows. A number of commercially available probiotic strains were inoculated into the milk drink and then treated at a pressure suitable for controlling contamination by yeasts and moulds, specifically 350 MPa for 5 minutes. The results of the selection method are shown in Table 1. Neither of the *Lactobacillus* strains evaluated survived the pressure treatment in sufficient numbers. However two of the *Bifidobacterium* strains sold under the trade marks "Bb12" (Nestle) and "Wisby 420" (Wisby), are not appreciably inactivated by the test-pressure treatment.

20

Table 1: Method of selecting a suitable probiotic strain

Species	Strains	Count (cfu/mL)		> 10^5 cfu/mL
		Before treatment	After treatment	
<i>L. casei</i>	shirota - Yakult	4.90×10^7	2.40×10^1	no
<i>L. johnsonii</i>	Lc-1- Nestle	7.80×10^7	9.00×10^1	no
<i>Bifidobacterium</i>	536 - Morinaga	1.30×10^8	9.00×10^1	no
<i>Bifidobacterium</i>	Bb12 - Nestle	6.20×10^7	5.20×10^7	yes
<i>Bifidobacterium</i>	420 - Wisby	5.70×10^7	3.30×10^7	yes

Example 18: Method of strain selection

Potential probiotic strains were screened for suitability in the production of a pressure-treated probiotic orange juice as follows. Two commercially available probiotic strains were inoculated into the juice (pH 4.0) and then treated at a pressure suitable for controlling contamination by spoilage bacteria, specifically 600 MPa for 5 minutes. The results of the selection method are shown in Table 2. Under these conditions both *Bifidobacterium* strains survive the pressure-treatment in sufficient numbers for delivery in a pressure treated orange juice.

Table 2: Method of selecting a suitable probiotic strain

Species	Strains	Count(cfu/mL)		> 10 ⁵ cfu/mL
		Before treatment	After treatment	
<i>Bifidobacterium</i>	Bb12 - Nestle	1.1 x 10 ⁸	4.1 x 10 ⁶	yes
<i>Bifidobacterium</i>	420 - Wisby	7.0 x 10 ⁷	1.2 x 10 ⁶	yes

Example 19: A low pH sports drink with a live probiotic culture

A whey protein isolate sport drink (4.8% whey protein isolate, protein, 0% fat and 10% carbohydrate), with a pH of 3.4, was inoculated with *Bifidobacterium lactis* strain HN019 at 9.2x10⁷ cfu/mL (RCA agar, Table 5) and intentionally contaminated by spiking at 4.5x10⁵ cfu/mL with pink yeast. The contaminated probiotic low pH sport drink was then treated at a pressure of 400 MPa for 5 minutes at 15 °C. This process produced a low pH sport drink with no detectable contaminating Pink yeast (> 4.6-log inactivation) while retaining a probiotic culture count of 2.7x10⁷ cfu/mL (RCA agar, Table 5).

Example 20: A nutrition shake with a live probiotic culture

A Myoplex French vanilla nutrition shake made by EAS, USA, (20 g protein (whey and soy protein isolates, 4.6 g fat and 20 g carbohydrate per 330mL serving). was inoculated with *Bifidobacterium lactis* strain HN019 at 5.5×10^7 cfu/mL and intentionally contaminated by spiking at 5.4×10^5 cfu/mL with Pink yeast. The contaminated probiotic neutral pH sport drink was than treated at a pressure of 500 MPa for 5 minutes at 15 °C. This process produced a neutral pH sport drink with no detectable contaminating Pink yeast (> 4.7 -log inactivation) while retaining a probiotic culture count of 4.8×10^7 cfu/mL.

Example 21 : A yoghurt containing a viable culture

A yoghurt milk made up with 7.0% SMP and 7.5% WMP, was heated to 55°C and homogenised at 150/50 bar. The homogenised milk was then heated to 90°C in a steam-heated water bath and held at that temperature for 10 minutes. After rapid cooling to 42°C, the yoghurt milk was inoculated with 1% of a starter culture sold by Rhodia under the trade mark "MY900", and fermented at 42°C to a pH of 4.1, whereupon it was cooled to 4°C. The yoghurt was pressure-treated at 400 MPa for 5 minutes (A treatment selected based on the inactivation of contaminating yeasts and moulds, as in earlier Examples). The treated yoghurt had a viable culture of 7.9×10^7 cfu/g (M17 agar, Table 6).

Example 22 : A yoghurt containing a viable culture

A yoghurt milk, made up with 12.0% whole milk powder (WMP) and 1.7% of whey protein concentrate was heated to 90°C in a steam-heated water bath and held at that temperature for 10 minutes. After cooling to 42°C, the yoghurt milk was inoculated with a starter culture sold by Rhodia under the trade mark "MY900" (commercial culture), and fermented at 42°C for 6 hours, to a pH of 4.4, Whereupon it was cooled to 4°C. The contaminated yoghurt was then pressure-treated at 450 MPa for about 5 minutes at 15°C. Based upon previous examples, this treatment is sufficient to inactivate typical dairy yeasts and moulds. This process produced a product with a starter culture count of 3.0×10^7 cfu/mL (M17 agar, Table 5).

Example 23 : A yoghurt with a viable culture

A yoghurt milk made up of 7.0% skim milk powder and 7.5% whole milk powder was heated to 55°C and homogenised at 150/50 bar. The homogenised milk was then heated to 90°C, held for 10 mins and rapidly cooled to 42°C. The milk was then inoculated with a Rhodia MY900 culture, fermented at 42°C to a pH of 4.5 and cooled to 4°C. Two samples of the resulting yoghurt were pressure treated at 350 MPa and 450 MPa for 1 minute and refrigerated. After 95 days of storage at 4°C, the live culture and contaminating mould counts were compared with an untreated control sample. The results are shown in Figure 2. The untreated and 350 MPa-treated samples were contaminated by more than one million colony-forming units of mould per gram, but the 450 MPa-treated sample had no detectable mould. In all three cases, the live culture count exceeded ten million colony-forming units per gram at 95 days.

Example 24 : A pressure yoghurt containing a viable culture

A yoghurt was made as in Example 24, and pressure treated to 430 MPa for about one second. After 15 days storage at 20°C, the level of live culture and contaminating moulds were compared with an untreated control sample. The results shown in Figure 3 show that after storage, the untreated control was contaminated by $>10^4$ cfu/mL of mould, but the pressure-treated yoghurt has no detectable mould. In both cases, the live culture exceeded 10^7 cfu/mL at 15 days.

Example 25 : A pressure treated yoghurt containing a viable culture

25

A yoghurt was made as in Example 24, intentionally spiked with $>10^5$ cfu/g of a penicillium mould, and pressure treated to the pressures 360 MPa, 400 MPa, 430 MPa, 460 MPa, 480 MPa and 500 MPa for about one second. The counts of live culture, moulds, and the mould log-reduction are given in Table 3, with an untreated control yoghurt for comparison. All pressure treatments produced yoghurts with $>9 \times 10^8$ cfu/g of live culture. The 360 MPa and 400 MPa pressure treatments did not significantly reduce

the level of mould contamination. The 430 MPa treatment produced a 2.4-log cycle reduction in mould counts, whereas the 460 MPa, 480 MPa and 500 MPa treatments produced yoghurts with no detectable moulds (>4.8 log-cycle reduction).

5 **Table 3**

Pressure treatment	Culture count (cfu/g)	Mould count (cfu/mL)	Mould reduction (cfu/mL)
None- control	1.2×10^9	7.3×10^5	0
360 MPa	1.1×10^9	$>1.5 \times 10^6$	(+0.3)
400 MPa	1.0×10^9	2.5×10^5	0.1
430 MPa	6.1×10^8	3.3×10^3	2.4
460 MPa	1.0×10^9	none detected (<10)	>4.8
480 MPa	1.1×10^9	none detected (<10)	>4.8
500 MPa	1.0×10^9	none detected (<10)	>4.8

Example 26 : A cultured beverage containing a live probiotic culture.

10

10% reconstituted skim milk in water was fermented with *Bifidobacterium lactis*, and mixed into a sucrose, carboxymethylcellulose mixture, pH-adjusted to 4.0 with phosphoric acid and homogenised at 200 bar, to produce a probiotic-containing beverage of 3% milk solids, 8% sugar and 0.4% carboxymethylcellulose. The beverage was pressure treated at 600 MPa for 1 second and the level of microbial contamination compared to an untreated control beverage. The pressure-treated and control-untreated beverages both contained live culture $> 10^8$ cfu/g, but the untreated beverage also had an aerobic plate count of 30 cfu/g and a yeast/mould count of 20 cfu/g. The pressure-treated beverage had no contaminants detected by those two methods.

20

Example 27 : Yoghurts made to three different pH values

Three yoghurts were made by fermenting 7.0% skim milk powder and 7.5% whole milk powder reconstituted in water with a culture of *Streptococcus thermophilus* (St10) and Lactobacillus bulgaricus (Lb1) to three different final pH values, 4.3, 4.1 and 3.9.

The three yoghurts were then pressure-treated at 350 MPa and 450 MPa for 5 minutes, and the surviving culture enumerated on M17 agar in each case, and shown in Figure 4.

The yoghurt at the higher pH (pH = 4.3) had live culture counts $>10^7$ cfu/mL after both treatments.

10 The yoghurt at the intermediate pH (pH = 4.1) had a live culture count $>10^7$ cfu/mL after the 350 MPa treatment, but not after the 450 MPa treatment

The yoghurt at the lower pH (pH = 3.9) had a live culture count $<10^7$ cfu/mL after both treatments.

15 **Example 28 : A yoghurt pressure treated for three different treatment times.**

10% reconstituted skim milk powder was inoculated with Lb-5033 and St-10 and fermented to a pH of 4.4 at 37°C. The resulting cultured milk was then challenged with a mixture of pink yeast and penicillium mould and pressure-treated at 350 MPa for either 5,

20 10 or 15mins. The cultured milk treated for 5 mins had a live culture count of 2.8×10^7 cfu/g (on M17 agar), and a 4.1 log-cycle reduction of contaminating yeasts and moulds,

The cultured milk treated for 10mins had a live culture count of only 5.7×10^5 cfu/g (M17 agar) with no detectable contaminating yeasts and moulds. The cultured milk

25 treated for 15mins had a live culture count of only 1.9×10^5 cfu/g (M17 agar) with no detectable contaminating yeasts and moulds.

Example 29 : Pressure treatment of commercial yoghurt cultures

30 Yoghurt milk was prepared by reconstituting SMP at 10% solids. The milk was heated to 55°C and homogenised at 150/50 bar, and then heated to 90°C for 10mins and cooled. The milk was then divided up into equal aliquots and each separately inoculated with a

commercial starter from the list in Table 4. The milks were incubated for 16 hours at 40°C or 38°C. After incubation, the yoghurts were cooled in an icebath and the pH measured, and adjusted to 4.4 as necessary. The yoghurts were each treated at a pressure of 400 MPa for 5 mins, and had a culture count of $>10^7$ cfu/g (on M17 agar).

5

Table 4

Culture	Supplier	400 MPa – 5min (count cfu/mL)
St-B-01	Chr. Hansen	2.0×10^3
YC-X11	Chr. Hansen	1.8×10^8
YC-180	Chr. Hansen	2.5×10^5
YC-380	Chr. Hansen	4.3×10^4
ABT 3	Chr. Hansen	6.0×10^7
ABT 4	Chr. Hansen	2.0×10^7
V1	Danisco	1.0×10^7
Yo Mix VW	Danisco	2.0×10^8
MSK Mix AB N 1-45	Danisco	2.3×10^8
Bf420	Danisco	3.6×10^5 *
MY 105	Rhodia	2.7×10^7
MYE 95	Rhodia	5.5×10^7
MY Bio 6	Rhodia	8.0×10^7
TA 060	Rhodia	1.3×10^8
LH 100	Rhodia	6.4×10^7 *
MY 900	Rhodia	2.2×10^8

*enumerated on MRS agar

10

Enumeration of culture

An appropriate amount of sample was added to diluent to produce a 10^{-1} dilution. Then, subsequent dilutions were prepared serially to plate out a range from 10^{-1} to 10^{-7} for each sample. The dilutions were then plated onto appropriate agars for each species (Table 5).

15

The agars were selected for efficient recovery of the test species, and prepared according to the manufacturer's instructions. The species and corresponding method used are summarised in Table 5.

Table 5: A Summary of the Methods used for Microbiological Enumeration

Culture	Agar used	Supplier	Summary of conditions
<i>S. thermophilus</i>	M17	Difco (Cockeysville, USA)	37°C, 48h, aerobic
<i>Lb. delbruekii</i> <i>subsp bulgaricus</i> <i>L. helveticus</i>	MRS	Difco	37°C, 48h, Anaerobic
Yeasts and moulds	YGCA	Merck Darmstadt, Germany	25°C, 5 days
<i>Bifidobacterium</i>	RCA	Merck	37°C, 3 days, Anaerobic
<i>L. rhamnosus</i> <i>L. johnsonii</i> <i>L. para-casei</i> <i>L. casei</i>	MRS	Difco	37°C, 3 days, Anaerobic
* <i>Bifidobacterium</i> differential	RBA	Made at Fonterra Research Centre	37°C, 3 days Anaerobic

RBA agar reference: Raffinose-Bifidobacterium (RB) agar, a new selective medium for Bifidobacteria', R. Hartemink et al, Journal of Microbial Methods 27 (1996) 33-43

Table 6 : A summary of the yeast and mould contaminants used in the examples

Strain	Origin	Source		Description
<i>Yarrowia lipolytica</i>	Yeast isolate from dairy product	FRC* Group collection	Starter culture	Contaminant from a dairy product
Unknown	'Pink yeast' Isolated from a dairy product	FRC Group collection	Starter culture	Contaminant from dairy product
<i>Candida famata</i>	Yeast isolate from dairy product	FRC Group collection	Starter culture	Contaminant from a dairy product
<i>Candida parapsilosis</i>	Yeast isolate from dairy product	FRC Group collection	Starter culture	Contaminant from a dairy product
<i>Debramyces Hanseii</i>	Yeast isolate from dairy product	FRC Group collection	Starter culture	Contaminant from a dairy product
<i>Geotrichum candidum</i>	Mould isolate from dairy product	FRC Group collection	Starter culture	Contaminant from a dairy product
<i>Cladosporium</i>	Mould isolate from dairy product	FRC Group collection	Starter culture	Contaminant from a dairy product
<i>Penicillium</i>	Mould isolate from dairy product	FRC Group collection	Starter culture	Contaminant from a dairy product

* Fonterra Research Centre, Palmerston North, New Zealand.

The above describes some preferred embodiments of the present invention and indicates several possible modifications but it will be appreciated by those skilled in the art that other modifications can be made without departing from the scope of the invention.

5 REFERENCES

1. E. Caplice and G.F. Fitzgerald, *International Journal of Food Microbiology*. 50 (1999) 131.
2. W.H. Holzapfel, R. Geisen, Schillinger, *International Journal of Food Microbiology*. 24 (1995) 343.
- 10 3. Y.-K. Lee and S. Salminen, *Trends in Food Science and Technology*, 6 (1995) 241.
4. K. Buchanan and M. Patterson, *The Chemical Engineer*, January 2002, 20.

What we claim is:

1. A method of treating a food comprising the following steps:
 - selecting a food comprising at least one strain of a culture, said strain
5 capable of surviving a pressure treatment at a predetermined pressure and pH, and
 - subjecting the food to a treatment pressure at or below the predetermined pressure, wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora.
- 10 2. A method according to claim 1 wherein the treatment pressure is at least 350MPa.
3. A method according to claim 2 wherein the treatment pressure is at least 400MPa.
- 15 4. A method according to any one of the preceding claims wherein the food is at a pH of between 3.0 and 8.0 when subjected to the treatment pressure.
5. A method according to claim 4 wherein the pH is between 3.6 and 4.8.
- 20 6. A method according to claim 5 wherein the pH is between 4.0 and 4.6.
7. A method according to any one of the preceding claims wherein the food is a cultured dairy product.
- 25 8. A method according to claim 7 wherein the cultured dairy product is yoghurt.
9. A method according to any one of claims 1 to 6 wherein the food is selected from a yoghurt drink, dairy dessert, cottage cheese, cream cheese and cultured beverages.

30

10. A method according to any one of the preceding claims wherein the strain of culture is selected from:

- i) *Lactobacillus acidophilus*
 - ii) *Bifidobacterium lactis*
 - 5 - iii) *Streptococcus thermophilus*;
 - iv) *Lactobacillus helveticus*;
 - v) *Lactobacillus delbrukeii subsp bulgaricus*;
- or any combination thereof.

10 11. A method of treating a food, comprising the steps:

- selecting a food comprising at least one strain of a culture, said strain being a probiotic strain capable of surviving a pressure treatment at a predetermined pressure and pH, and
- 15 - subjecting the food to a treatment pressure at or below the predetermined pressure, wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora.

12. A method according to claim 11 wherein the probiotic strain is *Bifidobacterium*.

20 13. A method according to claim 12 wherein the probiotic strain is *Bifidobacterium lactis*.

14. A method according to claim 13 wherein the probiotic strain is *Bifidobacterium lactis* HN019 AGAL deposit number NM 97/09513 dated 18 August 1997.

25

15. A method according to claim 11 wherein the probiotic strain is *Lactobacillus*.

16. A method according to claim 15 wherein the probiotic strain is *Lactobacillus acidophilus*.

30

17. A method according to claim 16 wherein the probiotic is *Lactobacillus acidophilus* HN017 AGAL deposit number NM 97/09515 dated 18 August 1997.
- 5 18. A method according to any one of claims 11 to 17 wherein the treatment pressure is at least 350MPa.
19. A method according to claim 18 wherein the treatment pressure is at least 400MPa.
- 10 20. A method according to claim 19 wherein the treatment pressure is at least 500MPa.
- 15 21. A method according to any one of claims 11 to 20 wherein the food is at a pH of between 3.0 and 4.6 when subjected to the treatment pressure.
22. A method according to any one of claims 11 to 21 wherein the food is selected from a yoghurt, a cultured dairy product, a beverage, a fruit juice or a vegetable juice.
- 20 23. A method of treating a food comprising the following steps:
- selecting a food comprising at least one strain of a protective culture, said strain capable of surviving a pressure treatment at a predetermined pressure and pH, and
 - 25 subjecting the food to a treatment pressure at or below the predetermined pressure, wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora.
- 30 24. The use of at least one bacterial strain in a food to be subjected to a treatment pressure at a predetermined pressure wherein the treatment pressure reduces, delays, prevents or eliminates growth of spoilage microflora, and the bacterial strain survives, said bacterial strain being selected from:

- i) *Lactobacillus acidophilus* HN017 AGAL deposit number NM97/09515 dated 18 August 1997;
 - ii) *Bifidobacterium lactis* HN019 AGAL deposit number NM97/09513 dated 18 August 1997;
 - 5 - iii) *Streptococcus thermophilus*;
 - iv) *Lactobacillus helveticus*;
 - v) *Lactobacillus delbrueckii subsp bulgaricus*;
 - vi) *Lactobacillus acidophilus*;
 - vii) *Bifidobacterium lactis*;
 - 10 or any combination thereof.
25. A method of treating a food comprising the following steps:
- selecting a food comprising *Lactobacillus acidophilus* HN017 AGAL deposit number NM97/09515 dated 18 August 1997; and
 - 15 - subjecting the food to a treatment pressure of between 350MPa and 600MPa, at a pH of between about 3.0 and about 8.0.
26. A method of treating a food comprising the following steps:
- selecting a food comprising *Bifidobacterium lactis* HN019 AGAL deposit
 - 20 number NM97/09513 dated 18 August 1997; and
 - subjecting the food to a treatment pressure of between 350MPa and 600MPa, at a pH of between about 3.0 and about 8.0.
27. A method according to any one of the preceding claims wherein the food is
- 25 subjected to the treatment pressure for less than 10 minutes.
28. A method according to claim 27 wherein the food is subjected to the treatment pressure for about 5 minutes.
- 30 29. A method according to claim 27 wherein the food is subjected to the treatment pressure less than 5 minutes.

30. A method according to claim 29 wherein the food is subjected to the treatment pressure for about 1 minute.
- 5 31. A method according to claim 29 wherein the food is subjected to the treatment pressure for less than 1 minute.
32. A method according to claim 31 wherein the food is subjected to the treatment pressure for less than 30 seconds.
- 10 33. A method according to claim 32 wherein the food is subjected to the treatment pressure for less than 5 seconds.
- 15 34. A method according to claim 33 wherein the food is subjected to the treatment pressure for about 1 second.
35. A method according to any one of the preceding claims wherein the food is subjected to the treatment pressure at a temperature between about 0 degrees Celsius and 40 degrees Celsius.
- 20 36. A method according to claim 35 wherein the food is subjected to the treatment pressure at a temperature between about 0 degrees Celsius and 20 degree Celsius.
37. A food prepared by method according to any one of the preceding claims.
- 25 38. A food according to claim 37 wherein the food is selected from a yoghurt, a cultured dairy product, a beverage or a fruit or vegetable juice.
- 30 39. A cultured dairy product having a pH of at least 4.0 and a viable culture of at least one hundred thousand colony-forming units per gram following a pressure treatment of at least 400 MPa.

40. A cultured dairy product with a pH of at least 4.0 having a viable culture of at least one hundred thousand colony-forming units per gram following a pressure treatment of at least 450 MPa.
- 5
41. A cultured dairy product with a pH of at least 4.0 having a viable culture of at least one hundred thousand colony-forming units per gram following a pressure treatment of at least 500 MPa.
- 10 42. A yoghurt or yoghurt drink with a pH of at least 4.0 having a viable culture of at least one hundred thousand colony-forming units per gram following a pressure treatment of at least 600MPa.
- 15 43. A food or beverage having a viable culture count of at least one hundred thousand colony-forming units per gram of at least one strain of a probiotic bacteria following a pressure treatment of at least 400 MPa for less than 10 mins.
- 20 44. A food or beverage having a viable culture count of at least one hundred thousand colony-forming units per gram of at least one strain of a probiotic bacteria following a pressure treatment of at least 450 MPa for less than 10 mins.
45. A method according to any one of claims 1 to 36 wherein the food has been packaged prior to being subjected to the treatment pressure.
- 25 46. Food made by the method according to any one of claims 1 to 36 wherein the spoilage organisms are inhibited for an extended period of time during storage, said extended period of time being longer than that achieved by an untreated food containing a strain of culture.
- 30 47. Food according to claim 46 wherein said storage is for at least 50 days at about 4 degrees Celsius.

48. Food according to claim 46 wherein said storage is for at least 90 days at about 4 degrees Celsius.
- 5 49. Food according to claim 46 wherein said storage is for at least 15 days at 20 degrees Celsius.

1/4

Figure 1a






Figure 1b



- 5 Effect of pressure treatment on survival of *Penicillium* mould (solid symbols) and Rhodia MY900 yoghurt starter (open symbols).

2/4


Figure 2



5 Yoghurt treated at different pressures with an abundant viable culture, but exhibiting varying degrees of spoilage after 95 days, depending on treatment pressure.

3/4


Figure 3



- 5 Effect of 1 second of pressure treatment at 430MPa on viable culture and spoilage levels after 15 days.

4/4

Figure 4



Effect of pH on pressure treatment

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ03/00224

A. CLASSIFICATION OF SUBJECT MATTERInt. Cl. ⁷: A23L 3/015, 3/3571

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

SEE ELECTRONIC DATA BASES

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SEE ELECTRONIC DATABASES

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPIDS, CA, FSTA (Pressure, culture, probiotic, bacteria, live, dairy, milk, yoghurt, juice, cheese, heat resistant).

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GB 2367997A. (KU Leuven Research and Development) 24 April 2002.	1-49
A	Ulmer, H. M. et al. 2000. "Effects of high pressure on survival and metabolic activity of <i>Lactobacillus plantarum</i> TMW1.460". Applied and Environmental Microbiology. 66 (9) pp 3966-3973.	1-49
A	Calik, H. et al. 2002. "Effect of high pressure processing on <i>Vibrio parahaemolyticus</i> strains in pure cultures and Pacific oysters". Journal of food science Vol: 67, 4, Pp 1506-1510.	1-49

☐

Further documents are listed in the continuation of Box C

☒

See patent family annex

* Special categories of cited documents:			
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

18 December 2003

Date of mailing of the international search report

23 DEC 2003

Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
E-mail address: pct@ipaaustralia.gov.au
Facsimile No. (02) 6285 3929

Authorized officer

Terry Moore

Telephone No : (02) 6283 2632

INTERNATIONAL SEARCH REPORT

International application No.

PCT/NZ03/00224

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
GB	20001129	D0	0025415	GB	20020424	A	2367997
							END OF ANNEX